

February 1, 2008

On the path integration for the potential barrier $V_0 \cosh^{-2}(\omega x)$

L. Guechi

Département de Physique,

Faculté des Sciences, Université Mentouri,

Route d'Ain El Bey, Constantine, Algérie.

T. F. Hammann

Laboratoire de Mathématiques, Physique mathématique,

Faculté des Sciences et Techniques,

Université de Haute Alsace,

4, rue des frères Lumière, F-68093 Mulhouse, France

Abstract

The propagator associated to the potential barrier $V = \cosh^{-2}(\omega x)$ is obtained by solving path integrals. The method of delta functionals based on canonical and other transformations is used to reduce the path integral for this potential into a path integral for the Morse potential problem. The dimensional extension technique is seen to be essential for performing the multiple integral representation of the propagator. The correctly normalized scattering wave functions and the scattering function are derived. To test the method employed, the free particle and the δ -function barrier are considered as limiting cases.

PACS 03.65.Ca-Formalism

PACS 03.65.Db-Functional analytical methods

typescript using Latex

I. Introduction

During the last twenty years, a number of nontrivial path integrals of nonrelativistic quantum mechanical problems have been solved^{1,2,3,4}. In particular, two wide classes for which the path integral solutions are related to the radial harmonic oscillator and the Pöschl-Teller path integral, are extensively studied. The first class called class of confluent potentials includes the radial harmonic oscillator, the Morse potential, and the Coulomb potential. The Pöschl-Teller and the modified Pöschl-Teller potentials, e.g. the Rosen-Morse, Manning-Rosen, Scarf-like, or the Hulthen potential belong to the class of hypergeometric potentials. However, most of discussions of these potentials by path integration were restricted to bound state problem. For the second class, when the extension to describe the scattering states was made, the approach applied is based on a Sommerfeld-Watson transformation which leads to a closed form expression for the Green's function^{5,6,7,8,9,10} or via the path integration over the $SU(1,1)$ manifold for the calculation of the propagator with bound and scattering states found simultaneously^{11,12,13,14,15}. Note that the noncompact group $SU(1,1)$ can be viewed as analytical continuation of the compact group $SU(2)$.

The purpose of this paper is to solve the problem of the potential barrier $V = \cosh^{-2}(\omega x)$ in the framework of Feynman's path integral. This potential is not a special case of the modified Pöschl-Teller potentials. It cannot be related to them by coordinate transformation and it has not a dynamical $SU(1,1)$ symmetry. Thus, neither the approach which consists in summing the spectral representations of the Green's function via a Sommerfeld-Watson transformation nor the path integration over the $SU(1,1)$ manifold can be applied to this potential. A treatment of this potential barrier has been presented recently in the Feynman approach through integration on the $SO(1,2)$ group¹⁶, but it involved several incorrect manipulations so that the amplitudes $T(k)$ and $R(k)$ of the transmitted and reflected beams obtained are not correct. We think that it is worthwhile to present another path integral approach to solve this potential barrier problem.

In sec.II, we first introduce an auxiliary dynamics by extension of the phase space in order to convert the path integral for the potential barrier into a path integral form analogous to the rigid rotator problem. Then, with the help of a point canonical transformation and the trick called "uncompleting the square"¹⁷, the propagator is connected to the path integral for the Morse

potential problem. In sec.III, the propagator is evaluated, from which we obtain the suitably normalized wave functions. The scattering function is determined from the asymptotic form of the wave functions in sec.IV. As a test of our calculation, two limiting cases are considered. The coefficients of the transmitted and reflected beams are found again for the free particle and for a particle subjected to the repulsive δ -function potential in sec.V. The section VI will be a conclusion.

II. Reduction of $\cosh^{-2}(\omega x)$ into a form of Morse Potential

The path integral in phase space for a particle of mass m subjected to the potential barrier defined by

$$V(x) = \frac{V_0}{\cosh^2(\omega x)}; V_0 > 0, \quad (1)$$

is written formally as follows:

$$K(x_f, x_i; T) = \int Dx Dp_x \exp \left\{ \frac{i}{\hbar} \int_0^T \left(p_x \dot{x} - \frac{p_x^2}{2m} - \frac{V_0}{\cosh^2(\omega x)} \right) dt \right\}. \quad (2)$$

In the time sliced representation, it is given by

$$K(x_f, x_i; T) = \lim_{N \rightarrow \infty} \int \prod_{j=1}^N dx_j \prod_{j=1}^{N+1} \frac{dp_{x_j}}{2\pi\hbar} \exp \left\{ \frac{i}{\hbar} \sum_{j=1}^{N+1} \left[p_{x_j} \Delta x_j - \left(\frac{p_{x_j}^2}{2m} + \frac{V_0}{\cosh^2(\omega x_j)} \right) \varepsilon \right] \right\}, \quad (3)$$

where we have adopted the standard notation:

$$\varepsilon = t_j - t_{j-1} = \frac{T}{N+1}, \Delta x_j = x_j - x_{j-1}, x_i = x(0) \text{ and } x_f = x(T) = x(t_{N+1}).$$

Let us first rescale (x, p_x) by $(\alpha = \omega x, p_\alpha = \frac{1}{\omega} p_x)$. Under this transformation, the propagator (3) becomes

$$K(x_f, x_i; T) = \omega \lim_{N \rightarrow \infty} \int \prod_{j=1}^N d\alpha_j \prod_{j=1}^{N+1} \frac{dp_{\alpha_j}}{2\pi\hbar} \exp \left\{ \frac{i}{\hbar} \sum_{j=1}^{N+1} \left[p_{\alpha_j} \Delta \alpha_j - \left(\frac{p_{\alpha_j}^2}{2m} + \frac{V_0}{\cosh^2(\alpha_j)} \right) \varepsilon \right] \right\}$$

$$\begin{aligned}
& - \left(\frac{p_{\alpha_j}^2}{2m} + \frac{V_0/\omega^2}{\cosh^2 \alpha_j} \right) \omega^2 \varepsilon \Big] \Big\} \\
= & \omega \int D\alpha Dp_\alpha \exp \left\{ \frac{i}{\hbar} \int_0^{\omega^2 T} \left[p_\alpha \dot{\alpha} \right. \right. \\
& \left. \left. - \left(\frac{p_\alpha^2}{2m} + \frac{V_0/\omega^2}{\cosh^2 \alpha} \right) \right] dt \right\}. \tag{4}
\end{aligned}$$

In analogy with the problem of a free particle moving on the two-dimensional sphere which is also known as the rigid rotator problem¹⁸, the path integration of the expression (4) can be simplified by introducing an additional dynamics by extension of the phase space with the help of the following path integral identity:

$$\begin{aligned}
& \exp \left[-\frac{i}{\hbar} \left(\int_0^{\omega^2 T} \frac{V_0/\omega^2}{\cosh^2 \alpha} dt \right) \right] \\
= & \int_{-\infty}^{\infty} d(\beta_f - \beta_i) \exp \left[-i \sqrt{\frac{2mV_0}{\hbar^2 \omega^2}} - \frac{1}{4} (\beta_f - \beta_i) \right] \lim_{N \rightarrow \infty} \int \prod_{j=1}^N d\beta_j \\
& \times \prod_{j=1}^{N+1} \frac{dp_{\beta_j}}{2\pi\hbar} \exp \left\{ \frac{i}{\hbar} \sum_{j=1}^{N+1} \left[p_{\beta_j} \triangle \beta_j - \frac{p_{\beta_j}^2 + \frac{\hbar^2}{4}}{2m \cosh^2 \alpha_j} \omega^2 \varepsilon \right] \right\} \\
= & \int_{-\infty}^{\infty} d(\beta_f - \beta_i) \exp \left[-i \sqrt{\frac{2mV_0}{\hbar^2 \omega^2}} - \frac{1}{4} (\beta_f - \beta_i) \right] \\
& \times \int D\beta Dp_\beta \exp \left\{ \frac{i}{\hbar} \int_0^{\omega^2 T} \left[p_\beta \dot{\beta} - \frac{p_\beta^2 + \frac{\hbar^2}{4}}{2m \cosh^2 \alpha} \right] dt \right\}. \tag{5}
\end{aligned}$$

Then, the propagator (4) becomes

$$\begin{aligned}
K(x_f, x_i; T) = & \omega \int_{-\infty}^{\infty} d(\beta_f - \beta_i) \exp \left[-i \sqrt{\frac{2mV_0}{\hbar^2 \omega^2}} - \frac{1}{4} (\beta_f - \beta_i) \right] \\
& \times \int D\alpha D\beta Dp_\alpha Dp_\beta \exp \left\{ \frac{i}{\hbar} \int_0^{\omega^2 T} \left[p_\alpha \dot{\alpha} + p_\beta \dot{\beta} \right. \right. \\
& \left. \left. - \left(\frac{p_\alpha^2}{2m} + \frac{p_\beta^2 + \frac{\hbar^2}{4}}{2m \cosh^2 \alpha} \right) \right] dt \right\}. \tag{6}
\end{aligned}$$

Our first task is to bring the (β, p_β) -dependent part of the above path integral into an appropriate form. To do this, we consider the kernel

$$\begin{aligned}
K(\beta_f, \beta_i; T) &= \int D\beta Dp_\beta \exp \left\{ \frac{i}{\hbar} \int_0^{\omega^2 T} \left[p_\beta \dot{\beta} - \frac{p_\beta^2 + \frac{\hbar^2}{4}}{2m \cosh^2 \alpha} \right] dt \right\} \\
&= \lim_{N \rightarrow \infty} \int \prod_{j=1}^N d\beta_j \prod_{j=1}^{N+1} \frac{dp_{\beta_j}}{2\pi\hbar} \\
&\quad \times \exp \left\{ \frac{i}{\hbar} \sum_{j=1}^{N+1} \left[p_{\beta_j} \Delta\beta_j - \frac{p_{\beta_j}^2 + \frac{\hbar^2}{4}}{2m \cosh^2 \alpha_j} \omega^2 \varepsilon \right] \right\}. \quad (7)
\end{aligned}$$

If we integrate this expression over the variables β_j , we obtain

$$\begin{aligned}
K(\beta_f, \beta_i; T) &= \lim_{N \rightarrow \infty} \int \frac{dp_{\beta_{N+1}}}{2\pi\hbar} \exp \left\{ \frac{i}{\hbar} [\beta_{N+1} p_{\beta_j} - \beta_0 p_{\beta_1}] \right\} \int \prod_{j=1}^N dp_{\beta_j} \\
&\quad \times \delta(p_{\beta_j} - p_{\beta_{j+1}}) \prod_{j=1}^{N+1} \exp \left(-\frac{i}{\hbar} \frac{p_{\beta_j}^2 + \frac{\hbar^2}{4}}{2m \cosh^2 \alpha_j} \omega^2 \varepsilon \right). \quad (8)
\end{aligned}$$

The p_{β_j} integrations thereafter give $p_{\beta_1} = p_{\beta_2} = \dots = p_{\beta_{N+1}} = \hbar\lambda$. Hence we have

$$K(\beta_f, \beta_i; T) = \int_{-\infty}^{\infty} \frac{d\lambda}{2\pi} \exp [i\lambda (\beta_f - \beta_i)] \exp \left(-\frac{i}{\hbar} \int_0^{\omega^2 T} \frac{\hbar^2}{2m \cosh^2 \alpha} \lambda^2 + \frac{1}{4} dt \right). \quad (9)$$

Substituting (9) into (6) yields the following expression for the propagator:

$$\begin{aligned}
K(x_f, x_i; T) &= \omega \int_{-\infty}^{\infty} d(\beta_f - \beta_i) \exp \left[-i\sqrt{\frac{2mV_0}{\hbar^2\omega^2} - \frac{1}{4}}(\beta_f - \beta_i) \right] \\
&\quad \times \int_{-\infty}^{\infty} \frac{d\lambda}{2\pi} \exp [i\lambda (\beta_f - \beta_i)] K(\alpha_f, \alpha_i; T), \quad (10)
\end{aligned}$$

where the (α, p_α) -part is given by

$$K(\alpha_f, \alpha_i; T) = \int D\alpha Dp_\alpha \exp \left[\frac{i}{\hbar} \int_0^{\omega^2 T} \left(p_\alpha \dot{\alpha} - \frac{p_\alpha^2}{2m} - \frac{\hbar^2}{2m \cosh^2 \alpha} \lambda^2 + \frac{1}{4} \right) dt \right]. \quad (11)$$

The calculation of this path integral is of course not at all straightforward. We will now follow Ref.¹⁷ to convert our propagator into the Morse potential problem by utilizing a set of tricks. We start by performing the point canonical transformation defined by

$$X = \sinh \alpha, P = \cosh \alpha p_\alpha, \quad (12)$$

which introduces the effective potential^{17,19}

$$V_e = \frac{\hbar^2}{8m} \left(3 - \frac{1}{1 + X^2} \right). \quad (13)$$

The transformed path integral takes the form

$$\begin{aligned} K(\alpha_f, \alpha_i; T) = & \left[(1 + X_f^2) (1 + X_i^2) \right]^{\frac{1}{4}} \int DX DP \exp \left\{ \frac{i}{\hbar} \int_0^{\omega^2 T} \left[P \dot{X} \right. \right. \\ & \left. \left. - \frac{P^2}{2m} (1 + X^2) - \frac{\hbar^2}{2m} \frac{\lambda^2}{1 + X^2} - \frac{3\hbar^2}{8m} \right] dt \right\}. \end{aligned} \quad (14)$$

One can get rid of the $(1 + X^2)^{-1}$ potential term, by changing the variable P into $\tilde{P}$ via

$$P = \tilde{P} - \frac{i\hbar\lambda}{1 + X^2}. \quad (15)$$

This lets appear a total X derivative which can easily be integrated out. Then, the path integral (14) may be rewritten

$$K(\alpha_f, \alpha_i; T) = \left[(1 + X_f^2) (1 + X_i^2) \right]^{\frac{1}{4}} \exp \left[\lambda (\arctan X) \Big|_{X_i}^{X_f} \right] K(X_f, X_i; T), \quad (16)$$

where

$$\begin{aligned} K(X_f, X_i; T) = & \int DX D\tilde{P} \exp \left\{ \frac{i}{\hbar} \int_0^{\omega^2 T} \left[\tilde{P} \dot{X} - \frac{\tilde{P}^2}{2m} (1 + X^2) \right. \right. \\ & \left. \left. + \frac{i\hbar\lambda}{m} \tilde{P} - \frac{3\hbar^2}{8m} \right] dt \right\}. \end{aligned} \quad (17)$$

Following Anderson et al¹⁷, the $\tilde{P}^2 X^2$ term can be absorbed using the trick called uncompleting the square, namely

$$\exp \left(-\frac{i}{\hbar} \int_0^{\omega^2 T} \frac{\tilde{P}^2}{2m} X^2 dt \right)$$

$$= \int_{-\infty}^{\infty} d\xi_f \int D\xi \exp \left[\frac{i}{\hbar} \int_0^{\omega^2 T} \left(\frac{m}{2} \dot{\xi}^2 - \dot{\xi} \tilde{P} X + \frac{3\hbar^2}{8m} \right) dt \right], \quad (18)$$

and (17) takes the form

$$K(X_f, X_i; T) = \int_{-\infty}^{\infty} d\xi_f \int D\xi \int DX D\tilde{P} \exp \left[\frac{i}{\hbar} \int_0^{\omega^2 T} \left(\frac{m}{2} \dot{\xi}^2 + \tilde{P} \dot{X} - \frac{\tilde{P}^2}{2m} - \dot{\xi} \tilde{P} X + \frac{i\hbar\lambda}{m} \tilde{P} \right) dt \right]. \quad (19)$$

By discretizing the $\tilde{P} \dot{X}$ term and performing the integrations on the N variables X_j , it appears N Dirac distributions which can formally be written as

$$\delta \left(\dot{\tilde{P}} + \dot{\xi} \tilde{P} \right). \quad (20)$$

Taking into account the boundary conditions $\tilde{P}(0) = P_0, \xi(0) = 0$ and $\xi(\omega^2 T) = \xi_f$, the solution of the differential equation in the argument of the Dirac distribution (20) is

$$\tilde{P}(t) = P_0 e^{-\xi(t)}. \quad (21)$$

The $\tilde{P}_j$ integrations thereafter may be performed and reduced to an ordinary integration over P_0 . It can also be shown that

$$\left(\frac{d\tilde{P}(\omega^2 T)}{dP_0} \right)^{\frac{1}{2}} = e^{-\frac{\xi_f}{2}}. \quad (22)$$

As a result, we have

$$K(X_f, X_i; T) = \int_{-\infty}^{\infty} d\xi_f \exp \left(-\frac{\xi_f}{2} \right) \int_{-\infty}^{\infty} \frac{dP_0}{2\pi\hbar} \times \exp \left[\frac{i}{\hbar} P_0 (e^{-\xi_f} X_f - X_i) \right] K_M(\xi_f, 0; T), \quad (23)$$

with

$$K_M(\xi_f, 0; T) = \int D\xi \exp \left[\frac{i}{\hbar} \int_0^{\omega^2 T} \left(\frac{m}{2} \dot{\xi}^2 - \frac{P_0^2}{2m} e^{-2\xi} + \frac{i\hbar\lambda}{m} P_0 e^{-\xi} \right) dt \right]. \quad (24)$$

This kernel describes the path integral for a particle to go from a position $\xi_i = 0$ at time $t_i = 0$ to ξ_f at time $t_f = \omega^2 T$, in the Morse-type potential

$$V(\xi) = \frac{P_0^2}{2m} \left(e^{-2\xi} - \frac{2i\hbar\lambda}{P_0} e^{-\xi} \right). \quad (25)$$

For $\frac{2i\hbar\mu}{P_0}$ real and positive, there exist treatments by path integral techniques^{1,2,19,20,21}, operator methods²² and algebraic approach²³. If one substitutes $A = \frac{P_0^2}{2m}$, $B = \frac{i\hbar\lambda}{m} P_0$, $a = 1$, $M = 4m$, $s = \frac{2m}{P_0} S$ in Ref.²³, one obtains

$$\begin{aligned} K_M(\xi_f, 0; T) &= \frac{2m}{\hbar} \int_0^\infty \frac{e^{-2\lambda S}}{i \sin(S)} dS \int_{-\infty}^\infty \frac{dE}{2\pi\hbar} \exp\left(-\frac{i}{\hbar} E \omega^2 T\right) \\ &\quad \times \exp\left[\frac{i}{\hbar} P_0 (e^{-\xi_f} + 1) \cot(S)\right] \\ &\quad \times I_{2\sqrt{-2mE/\hbar^2}} \left(\frac{2P_0 e^{-\frac{\xi_f}{2}}}{i\hbar \sin(S)} \right). \end{aligned} \quad (26)$$

With the results (26), (23) and (16) the full propagator (10) for the potential under consideration may be put into the form

$$\begin{aligned} K(x_f, x_i; T) &= \frac{2m\omega}{\hbar} [(1 + X_f^2)(1 + X_i^2)]^{\frac{1}{4}} \exp\left[\lambda(\arctan X_f - \arctan X_i)\right] \\ &\quad \times \int_{-\infty}^\infty d(\beta_f - \beta_i) \exp\left[-i\sqrt{\frac{2mV_0}{\hbar^2\omega^2}} - \frac{1}{4}(\beta_f - \beta_i)\right] \\ &\quad \times \int_{-\infty}^\infty \frac{d\lambda}{2\pi} \exp[i\lambda(\beta_f - \beta_i)] \int_{-\infty}^\infty d\xi_f \exp\left(-\frac{\xi_f}{2}\right) \int_{-\infty}^\infty \frac{dP_0}{2\pi\hbar} \\ &\quad \times \exp\left[\frac{i}{\hbar} P_0 (e^{-\xi_f} X_f - X_i)\right] \int_0^\infty \frac{e^{-2\lambda S}}{i \sin(S)} dS \int_{-\infty}^\infty \frac{dE}{2\pi\hbar} \\ &\quad \times \exp\left(-\frac{i}{\hbar} E \omega^2 T\right) \exp\left[\frac{i}{\hbar} P_0 \cot(S) (e^{-\xi_f} + 1)\right] \\ &\quad \times I_{2\sqrt{-2mE/\hbar^2}} \left(\frac{2P_0 e^{-\frac{\xi_f}{2}}}{i\hbar \sin(S)} \right), \end{aligned} \quad (27)$$

where $X_f = \sinh \alpha_f$ and $X_i = \sinh \alpha_i$.

III. Evaluation of the propagator

In order to evaluate the multiple integral (27), we start by employing the integral representation for the modified Bessel function

$$-2\pi i I_\mu(-2iaz) = z^{2\mu} \int_{-\infty}^{\infty} du u^{-(1+2\mu)} \exp \left[-ia \left(u + \frac{z^2}{u} \right) \right], \quad (28)$$

valid for $\arg(a) = 0$ and $\text{Re}(2\mu) \succ -1$, changing variables $q = e^{-\xi_f}$, $\tilde{q} = q/u^2$ and $\tilde{P}_0 = P_0/\sin(S)$ and carrying out the integration over $\tilde{P}_0$. We then obtain

$$\begin{aligned} K(x_f, x_i; T) &= \frac{2m\omega}{\hbar} [(1 + X_f^2)(1 + X_i^2)]^{\frac{1}{4}} \exp \left[\lambda(\arctan X_f - \arctan X_i) \right] \\ &\times \int_{-\infty}^{\infty} d(\beta_f - \beta_i) \exp \left[-i \sqrt{\frac{2mV_0}{\hbar^2 \omega^2} - \frac{1}{4}} (\beta_f - \beta_i) \right] \\ &\times \int_{-\infty}^{\infty} \frac{d\lambda}{2\pi} \exp [i\lambda (\beta_f - \beta_i)] \int_{-\infty}^{\infty} \frac{dE}{2\pi\hbar} \\ &\times \exp \left(-\frac{i}{\hbar} E \omega^2 T \right) \int_0^{\infty} dS e^{-2\lambda S} \int_0^{\infty} \tilde{q}^{\mu-\frac{1}{2}} d\tilde{q} \\ &\times \int_{-\infty}^{\infty} \delta(\tilde{q}\mathfrak{B}u^2 - u(1 + \tilde{q}) + \mathfrak{A}) du, \end{aligned} \quad (29)$$

where we have set

$$\mathfrak{A} = \cos(S) + X_f \sin(S), \quad \mathfrak{B} = \cos(S) - X_i \sin(S), \quad (30)$$

and

$$\mu = \sqrt{-\frac{2mE}{\hbar^2}} = \pm \frac{ik}{\omega}. \quad (31)$$

Next, the integration over u yields

$$\int_{-\infty}^{\infty} \delta(\tilde{q}\mathfrak{B}u^2 - u(1 + \tilde{q}) + \mathfrak{A}) du = \frac{2}{\sqrt{\tilde{q}^2 + 2(1 - 2\mathfrak{A}\mathfrak{B})\tilde{q} + 1}}. \quad (32)$$

If we let

$$S = \tilde{S} + \frac{1}{2} (\arctan X_f - \arctan X_i), \quad \Delta = -(\arctan X_f + \arctan X_i), \quad (33)$$

and noticing that

$$\cos(S) \mp X \sin(S) = \sqrt{1 + X^2} \cos(S \pm \arctan X), \quad (34)$$

the factor $(1 - 2\mathfrak{A}\mathfrak{B})$ is amenable to the form

$$(1 - 2\mathfrak{A}\mathfrak{B}) = \sinh \alpha_f \sinh \alpha_i - \cosh \alpha_f \cosh \alpha_i \cos(2\tilde{S}). \quad (35)$$

The integrations over the variables $\tilde{q}$, λ and $\tilde{S}$ can be easily performed using the integral (see formula 3.252.11 p.297 in²⁴)

$$\begin{aligned} & \int_0^\infty (1 + 2\beta x + x^2)^{\mu - \frac{1}{2}} x^{-\nu - 1} dx \\ &= 2^{-\mu} (\beta^2 - 1)^{\frac{\mu}{2}} \Gamma(1 - \mu) B(\nu - 2\mu + 1, -\nu) P_{\nu - \mu}^\mu(\beta), \end{aligned} \quad (36)$$

valid for $Re(\nu) < 0$, $Re(2\mu - \nu) < 1$, and $|\arg(\beta \pm 1)| < \pi$, where $B(x, y) = \frac{\Gamma(x)\Gamma(y)}{\Gamma(x+y)}$.

Taking into account the symmetry property of the general associated Legendre functions (see formula 8.731.5 p.1005 in²⁴)

$$P_{-\frac{1}{2} - i\lambda}^\mu(x) = P_{-\frac{1}{2} + i\lambda}^\mu(x), \quad (37)$$

and the well-known relation (see formula 8.334.2 p.937 in²⁴)

$$\Gamma\left(\frac{1}{2} - x\right)\Gamma\left(\frac{1}{2} + x\right) = \frac{\pi}{\cos \pi x}, \quad (38)$$

we obtain

$$\begin{aligned} K(x_f, x_i; T) &= \frac{\sqrt{\cosh \alpha_f \cosh \alpha_i}}{\pi \omega^2} \int_{-\infty}^\infty \frac{k dk}{\cosh\left(\frac{\pi k}{\omega}\right)} \exp\left(-\frac{i}{\hbar} \frac{\hbar^2 k^2}{2m} T\right) \\ &\quad \times \int_{-\infty}^\infty d(\beta_f - \beta_i) \exp\left[-i \sqrt{\frac{2mV_0}{\hbar^2 \omega^2} - \frac{1}{4}} (\beta_f - \beta_i)\right] \\ &\quad \times P_{-\frac{1}{2} + \frac{i k}{\omega}}(-Z), \end{aligned} \quad (39)$$

where

$$Z = \cosh \alpha_f \cosh \alpha_i \cosh(\beta_f - \beta_i) - \sinh \alpha_f \sinh \alpha_i. \quad (40)$$

The variables $\alpha_f, \alpha_i, \beta_f$, and β_i in (39) can be separated with the help of the addition theorem derived in Ref.¹⁶

$$P_{-\frac{1}{2}+\frac{ik}{\omega}}(-Z) = \frac{1}{4\pi} \sum_{\varepsilon=0}^1 \int_{-\infty}^{\infty} d\lambda e^{-i\lambda(\beta_f-\beta_i)} d_{\lambda,0}^{-\frac{1}{2}+\frac{ik}{\omega},\varepsilon}(\alpha_f+i\pi) \left(d_{\lambda,0}^{-\frac{1}{2}+\frac{ik}{\omega},\varepsilon}(\alpha_i) \right)^*, \quad (41)$$

where

$$d_{\lambda,0}^{-\frac{1}{2}+\frac{ik}{\omega},\varepsilon}(\alpha) = \frac{1}{2\pi} \int_{-\infty}^{\infty} d\beta \exp(-i\lambda\beta) [\cosh \beta \cosh \alpha + (-1)^\varepsilon \sinh \alpha]^{-\frac{1}{2}+\frac{ik}{\omega}}, \quad (42)$$

which may be evaluated by using the integral (see formula 8.713.3 p.1001 in²⁴)

$$P_\nu^{-\gamma}(z) = \sqrt{\frac{2}{\pi}} \frac{\Gamma(\gamma + \frac{1}{2}) (z^2 - 1)^{\frac{\gamma}{2}}}{\Gamma(\nu + \gamma + 1) \Gamma(\gamma - \nu)} \int_0^\infty \frac{\cosh[(\nu + \frac{1}{2})t] dt}{(z + \cosh t)^{\gamma + \frac{1}{2}}}, \quad (43)$$

valid for $Re(z) \succ -1, |\arg(z \pm 1)| \prec \pi, Re(\nu + \gamma) \succ -1, Re(\nu - \gamma) \succ 0$, and we have

$$d_{\lambda,0}^{-\frac{1}{2}+\frac{ik}{\omega},\varepsilon}(\alpha) = \frac{\Gamma(\frac{1}{2} + i\lambda - \frac{ik}{\omega}) \Gamma(\frac{1}{2} - i\lambda - \frac{ik}{\omega})}{\sqrt{2\pi} \Gamma(\frac{1}{2} - \frac{ik}{\omega})} \frac{e^{-\frac{\pi k}{2\omega}}}{\sqrt{\cosh \alpha}} P_\omega^{\frac{ik}{\omega}}(-\frac{1}{2} + i\lambda) ((-1)^\varepsilon \tanh \alpha). \quad (44)$$

Inserting (44) and (41) in (39) and integrating over the variable $(\beta_f - \beta_i)$, we arrive at

$$\begin{aligned} K(x_f, x_i; T) &= -\frac{1}{2\omega} \int_{-\infty}^{\infty} \frac{k dk}{|\sin[\pi(\nu - \frac{ik}{\omega})]|^2} e^{-\frac{\pi k}{\omega}} \sum_{\varepsilon=0}^1 \left(P_\omega^{\frac{ik}{\omega}}((-1)^\varepsilon \tanh \alpha_f) \right. \\ &\quad \times \left. P_\omega^{-\frac{ik}{\omega}}((-1)^\varepsilon \tanh \alpha_i) \right) \exp\left(-\frac{i}{\hbar} \frac{\hbar^2 k^2}{2m} T\right), \end{aligned} \quad (45)$$

where

$$\nu = -\frac{1}{2} + i\lambda = \frac{1}{2} \left(-1 + \sqrt{1 - \frac{8mV_0}{\hbar^2 \omega^2}} \right). \quad (46)$$

In order to find the wave functions, we set

$$\Psi_1(y) = P_\nu^{-\mu}(y), \quad \Psi_2(y) = P_\nu^{-\mu}(-y). \quad (47)$$

The parameter ν can be real or complex. In the complex case, we have

$$1 + \nu^* = -\nu. \quad (48)$$

Owing to (37), it is obvious that

$$\Psi_1^*(y) = P_\nu^\mu(y), \quad \Psi_2^*(y) = P_\nu^\mu(-y), \quad (49)$$

and thanks to the relations (see formulas 8.737.1 and 8.737.2 p.1006 in²⁴)

$$\begin{cases} P_\nu^{-\mu}(y) = \frac{\Gamma(\nu-\mu+1)}{\Gamma(\nu+\mu+1)} [\cos(\pi\mu)P_\nu^\mu(y) - \frac{2}{\pi}\sin(\pi\mu)Q_\nu^\mu(y)], \\ P_\nu^\mu(-y) = \cos[\pi(\nu+\mu)]P_\nu^\mu(y) - \frac{2}{\pi}\sin[\pi(\nu+\mu)]Q_\nu^\mu(y), \end{cases} \quad (50)$$

it is easy to make up the following equations

$$\begin{cases} \Psi_1(y) = a\Psi_1^*(y) + b\Psi_2^*(y), \\ \Psi_2^*(y) = a^*\Psi_2(y) + b^*\Psi_1(y), \end{cases} \quad (51)$$

where

$$a = \frac{\Gamma(\nu - \mu + 1) \sin(\pi\nu)}{\Gamma(\nu + \mu + 1) \sin[\pi(\nu + \mu)]}, \quad b = \frac{\Gamma(\nu - \mu + 1) \sin(\pi\mu)}{\Gamma(\nu + \mu + 1) \sin[\pi(\nu + \mu)]}. \quad (52)$$

Expression (45) of the propagator is then rewritten

$$\begin{aligned} K(x_f, x_i; T) = & -\frac{1}{2\omega} \int_{-\infty}^{\infty} \frac{k dk}{\left| \sin \left[\pi \left(\nu - \frac{ik}{\omega} \right) \right] \right|^2} e^{-\frac{\pi k}{\omega}} (\Psi_1(y_f) \Psi_1^*(y_i) \\ & + \Psi_2(y_f) \Psi_2^*(y_i)) \exp \left(-\frac{i}{\hbar} \frac{\hbar^2 k^2}{2m} T \right). \end{aligned} \quad (53)$$

Next by changing k into $(-k)$ in the interval of integration $]-\infty, 0]$, and substituting $\Psi_j(y)$ and $\Psi_j^*(y)$ by their expressions (51), we get

$$\begin{aligned} K(x_f, x_i; T) = & \frac{m}{2\hbar^2\omega} \int_0^{\infty} dE_k \frac{\sinh \left(\frac{\pi k}{\omega} \right)}{\left| \sin \left[\pi \left(\nu - \frac{ik}{\omega} \right) \right] \right|^2} \exp \left(-\frac{i}{\hbar} E_k T \right) \\ & \times \{ (|a|^2 + |b|^2) (\Psi_1(y_f) \Psi_1^*(y_i) + \Psi_2(y_f) \Psi_2^*(y_i)) \\ & + (ab^* + a^*b) (\Psi_1(y_f) \Psi_2^*(y_i) + \Psi_2(y_f) \Psi_1^*(y_i)) \}. \end{aligned} \quad (54)$$

Now it is easy to get assured that

$$|a|^2 + |b|^2 = 1, (ab^* + a^*b) = 0, \forall \nu. \quad (55)$$

Finally, the expression of the propagator is given by

$$\begin{aligned} K(x_f, x_i; T) = & \frac{m}{2\hbar^2\omega} \int_0^\infty dE_k \frac{\sinh\left(\frac{\pi k}{\omega}\right)}{\left|\sin\left[\pi\left(\nu - \frac{ik}{\omega}\right)\right]\right|^2} \\ & \times \{\Psi_1(y_f)\Psi_1^*(y_i) + \Psi_2(y_f)\Psi_2^*(y_i)\} \exp\left(-\frac{i}{\hbar}E_k T\right). \end{aligned} \quad (56)$$

The suitably normalized wave functions, corresponding to the energy $E_k = \frac{\hbar^2 k^2}{2m}$, are readily obtained to be

$$\Psi_{\leftarrow}(y) = \sqrt{\frac{m}{2\hbar^2\omega}} \frac{\left[\sinh\left(\frac{\pi k}{\omega}\right)\right]^{\frac{1}{2}}}{\left|\sin\left[\pi\left(\nu - \frac{ik}{\omega}\right)\right]\right|} P_{\nu}^{\frac{ik}{\omega}}(\pm \tanh \alpha), \quad (57)$$

where we have used the standard convention according to which the arrow in the wave function indicates the way in which the waves are propagating.

Taking into account the well-known link between the general associated Legendre functions and the hypergeometric functions (see formula 8.704 p.999 in²⁴)

$$P_{\nu}^{\mu}(x) = \frac{1}{\Gamma(1-\mu)} \left[\frac{1+x}{1-x}\right]^{\frac{\mu}{2}} F\left(-\nu, \nu+1; 1-\mu; \frac{1-x}{2}\right), \quad (58)$$

and using the relation (see formula 9.131.1 p.1043 in²⁴)

$$F(\alpha, \beta; \gamma; z) = (1-z)^{\gamma-\alpha-\beta} F(\gamma-\alpha, \gamma-\beta; \gamma; z), \quad (59)$$

the wave functions are written

$$\begin{aligned} \Psi_{\leftarrow}(y) = & \sqrt{\frac{m}{2\hbar^2\omega}} \frac{\left[\sinh\left(\frac{\pi k}{\omega}\right)\right]^{\frac{1}{2}}}{\left|\sin\left[\pi\left(\nu - \frac{ik}{\omega}\right)\right]\right|} \frac{1}{\Gamma(1-\frac{ik}{\omega})} \left[\frac{1-\tanh^2 \alpha}{4}\right]^{-\frac{ik}{2\omega}} \\ & \times F\left(1+\nu - \frac{ik}{\omega}, -\nu - \frac{ik}{\omega}; 1 - \frac{ik}{\omega}; \frac{1 \mp \tanh \alpha}{2}\right). \end{aligned} \quad (60)$$

IV. Scattering function

The sum of the reflection and transmission amplitudes $R(k)$ and $T(k)$ or else the scattering function $S(k)$ is independent of the way in which the waves are propagating. As an example, we can consider the case described by $\Psi_{\rightarrow}(y)$ in order to evaluate $S(k)$.

Using the identities for the hypergeometric functions (see formula 9.131.2 p.1043 in²⁴)

$$F\left(\alpha, \beta; \gamma; \frac{1-y}{2}\right) \rightarrow \begin{cases} 1, & y \rightarrow 1, \\ \frac{\Gamma(\gamma)\Gamma(\gamma-\alpha-\beta)}{\Gamma(\gamma-\alpha)\Gamma(\gamma-\beta)} + \left(\frac{1+y}{2}\right)^{\gamma-\alpha-\beta} \frac{\Gamma(\gamma)\Gamma(\alpha+\beta-\gamma)}{\Gamma(\alpha)\Gamma(\beta)}, & y \rightarrow -1, \end{cases} \quad (61)$$

we obtain

$$\begin{aligned} \Psi_{\rightarrow}(y) &\rightarrow \sqrt{\frac{m}{2\hbar^2\omega}} \frac{[\sinh(\frac{\pi k}{\omega})]^{\frac{1}{2}}}{|\sin[\pi(\nu - \frac{ik}{\omega})]|} \\ &\times \begin{cases} \frac{1}{\Gamma(1-\frac{ik}{\omega})} e^{ikx}, & x \rightarrow +\infty \\ \frac{\Gamma(-\frac{ik}{\omega})}{\Gamma(1+\nu-\frac{ik}{\omega})\Gamma(-\nu-\frac{ik}{\omega})} e^{ikx} + \frac{\Gamma(\frac{ik}{\omega})}{\Gamma(1+\nu)\Gamma(-\nu)} e^{-ikx}, & x \rightarrow -\infty. \end{cases} \end{aligned} \quad (62)$$

Identifying these asymptotic behaviors with the boundary conditions

$$\Psi_{\rightarrow}(y) \rightarrow \begin{cases} T(k)e^{ikx}, & x \rightarrow +\infty \\ e^{ikx} + R(k)e^{-ikx}, & x \rightarrow -\infty, \end{cases} \quad (63)$$

the amplitudes $T(k)$ and $R(k)$ of the transmitted and reflected beams are evaluated to be

$$\begin{cases} T(k) = \frac{\Gamma(1+\nu-\frac{ik}{\omega})\Gamma(-\nu-\frac{ik}{\omega})}{\Gamma(1-\frac{ik}{\omega})\Gamma(-\frac{ik}{\omega})}, \\ R(k) = \frac{\Gamma(1+\nu-\frac{ik}{\omega})\Gamma(-\nu-\frac{ik}{\omega})\Gamma(\frac{ik}{\omega})}{\Gamma(1+\nu)\Gamma(-\nu)\Gamma(-\frac{ik}{\omega})}, \end{cases} \quad (64)$$

from which the expression of the scattering function is

$$S(k) = T(k) + R(k) = \frac{\Gamma(\frac{ik}{\omega})\Gamma(-\nu-\frac{ik}{\omega})\cos[(\frac{\pi}{2})(\nu+\frac{ik}{\omega})]}{\Gamma(-\frac{ik}{\omega})\Gamma(-\nu+\frac{ik}{\omega})\cos[(\frac{\pi}{2})(\nu-\frac{ik}{\omega})]}, \quad (65)$$

and is clearly unitary

$$S^*(k) = S^{-1}(k). \quad (66)$$

Note that the analysis of the asymptotic behavior of $\Psi_{\leftarrow}(y)$ with the boundary conditions

$$\Psi_{\leftarrow}(y) \rightarrow \begin{cases} T(k)e^{-ikx}, & x \rightarrow -\infty \\ e^{-ikx} + R(k)e^{ikx}, & x \rightarrow +\infty, \end{cases} \quad (67)$$

leads to the same expression of the scattering function.

The exact expressions for $T(k)$ and $R(k)$ in Eqs. (64) differ significantly from the ones deduced from the solution¹⁶ obtained through path integration over $SO(1, 2)$.

V. Particular cases

We consider now two limiting cases.

A. The free particle

We have to check whether for $V_0 \rightarrow 0$, we obtain the well-known coefficients $T(k)$ and $R(k)$ relative to a free particle.

From Eq. (46), we notice that

$$\frac{\nu}{V_0 \rightarrow 0} \rightarrow 0. \quad (68)$$

Thus, we obtain from Eqs. (64):

$$\lim_{\nu \rightarrow 0} T(k) = 1, \quad \lim_{\nu \rightarrow 0} R(k) = 0. \quad (69)$$

These equations obviously prove that the particle is free.

B. The δ -function barrier

By letting $V_0 \rightarrow \infty$ and $\omega \rightarrow \infty$ such that $\frac{2V_0}{\omega} \rightarrow \frac{\hbar^2}{2m}g = \text{const.}$, we have

$$\int_{-\infty}^{\infty} \frac{V_0}{\cosh^2(\omega x)} dx = \frac{\hbar^2}{2m}g, \quad (70)$$

hence the potential (1) becomes

$$V(x) = \frac{\hbar^2}{2m}g\delta(x). \quad (71)$$

In this case

$$\lim_{(V_0 \rightarrow \infty, \omega \rightarrow \infty)} \nu \rightarrow -\frac{g}{2\omega}. \quad (72)$$

Thus,

$$\lim_{(V_0 \rightarrow \infty, \omega \rightarrow \infty)} T(k) = \frac{2k}{2k + ig}, \quad \lim_{(V_0 \rightarrow \infty, \omega \rightarrow \infty)} R(k) = \frac{-ig}{2k + ig}. \quad (73)$$

One can then easily see that if one substitutes $g = -\gamma$, one obtains the amplitudes of transmitted and reflected beams for the attractive δ -function barrier given in Ref²⁵.

VI. Conclusion

The main result of this article concerns the construction of the propagator relative to the potential barrier $V = \cosh^{-2}(\omega x)$. In carrying out the path integration, the method of delta functionals has enabled us to connect the potential barrier to the Morse potential problem. The evaluation of the multiple integral representation of the propagator is facilitated by using the technique of dimensional extension.

It seems to us that this method is a valuable alternative to the treatment of this potential by path integration over the $SO(1, 2)$ manifold.

We have also extracted the correctly normalized wave functions. The scattering function is obtained from their asymptotic behavior. Two limiting cases are considered. They constitute a consistency check for the correctness of our results.

References

- ¹D.C. Khandekar and S.V. Lawande, Phys. Rep. **137** (1986) 115; D.C. Khandekar, S.V. Lawande and K.V. Bhagwat, Path Integral Methods and Their Applications (World Scientific, Singapore, 1993).
- ²H. Kleinert, Path Integrals in Quantum Mechanics, Statistics and Polymer Physics (World Scientific, Singapore, 1990).
- ³A. Inomata, H. Kuratsuji and C.C. Gerry, Path Integrals and Coherent States of $SU(2)$ and $SU(1,1)$ (World Scientific, Singapore, 1992).
- ⁴C. Grosche and F. Steiner, Handbook of Feynman Path Integrals (Springer Tracts in Modern Physics, 1998).
- ⁵I.H. Duru and H. Kleinert, Fortschr. Phys. **30** (1982) 401.
- ⁶H. Kleinert and I. Mustapic, J. Math. Phys. **33** (1992) 643.
- ⁷L. Chetouani, L. Guechi, A. Lecheheb and T. F. Hammann, J. Math. Phys. **34** (1993) 1257.
- ⁸L. Chetouani, L. Guechi, A. Lecheheb and T.F. Hammann, Czech. J. Phys. **45** (1995) 699.
- ⁹L. Chetouani, L. Guechi, A. Lecheheb , T. F. Hammann and A. Messouber, Nuovo Cimento **B113** (1998) 81.
- ¹⁰C. Grosche, J. Phys. A: Math. Gen. **29** (1996) 365.
- ¹¹G. Junker and M. Böhm, Phys. Lett. **A117** (1986) 375.
- ¹²M. Böhm and G. Junker, J. Math. Phys. **28** (1987) 1978.
- ¹³A.O. Barut, A. Inomata and G. Junker, J. Phys. A: Math. Gen. **20**, (1987) 6271; **23**, (1990) 1179.
- ¹⁴T. Boudjedaa, L. Chetouani, L. Guechi and T. F. Hammann, J. Math. Phys. **32** (1991) 441.
- ¹⁵C. Grosche, J. Math. Phys. **32** (1991) 1984.

- ¹⁶H. Ahmedov and I.H. Duru, J. Phys. A: Math. Gen. **30** (1997) 173.
- ¹⁷A. Anderson and S.B. Anderson, Ann.Phys. (N.Y) **199** (1990) 155.
- ¹⁸D. Peak and A. Inomata, J. Math. Phys. **10** (1969) 1422; W. Langguth and A. Inomata, J. Math. Phys. **20** (1979) 499.
- ¹⁹N.K. Pak and I. Sökmen, Phys. Rev. **A 30** (1984) 1629.
- ²⁰P.Y. Cai, A. Inomata and R. Wilson, Phys. Lett. **A96** (1983) 117.
- ²¹I.H. Duru, Phys. Rev. **D28** (1983) 2689.
- ²²M.M. Nieto and L. M.Jr. Simmons, Phys. Rev. **D20** (1979) 2689.
- ²³L. Chetouani, L. Guechi and T.F. Hammann, Helv. Phys. Acta **65** (1992) 1069.
- ²⁴I.S. Gradshtein and I.M. Ryzhik, Tables of Integrals, Series and Products (Academic Press, NewYork 1965).
- ²⁵K. Gottfried, Quantum Mechanics, Vol. I : Fundamentals (W.A. Benjamin, Inc. London 1966).